

SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

- Operators must wear suitable protective clothing (coveralls) and suitable protective gloves when handling the concentrate.
- Operators must wear suitable protective gloves when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin or eyes immediately.

DO NOT BREATHE SPRAY.

WASH HANDS AND EXPOSED SKIN before meals and after work.

DO NOT APPLY via hand held equipment.

Consumer protection:

NOT TO BE USED ON FOOD CROPS.

Environmental protection:

Livestock must be kept out of treated areas for at least 7 days following treatment. IF RAGWORT IS PRESENT, FOLLOW THE GUIDANCE IN THE 'DIRECTIONS FOR USE'.

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with the LERAP requirement.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 metres of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 metre of the top of a ditch which is dry at the time of application. Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with HSE's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

LERAP scheme not applicable in the Republic of Ireland

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

Storage and disposal:

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

KEEP AWAY from food, drink and animal feeding stuffs.

KEEP OUT of reach of children.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

 **Dow AgroSciences**



Product Registration Number: MAPP 17772

An emulsifiable concentrate containing 150 g ae/litre (14.68% w/w) fluroxypyr (present as 216 g/litre of fluroxypyr-meptyl) and 150 g ae/litre (14.68% w/w) triclopyr (present as 209 g/litre triclopyr butyl).

A foliar acting herbicide for the control of DOCKS in ESTABLISHED GRASSLAND

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.

PROTECT FROM FROST.

2 Litres e

Dow AgroSciences Limited


CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE

Telephone: (01462) 457272 Fax: (01462) 426605

24 Hour Emergency Telephone Number: +44 (0) 1553 761 251

Trade-mark of The Dow Chemical Company ("Dow") or an Affiliated Company of Dow

This label is compliant with the CPA Voluntary Initiative Guidance (UK only)



Product Identifier according to Art. 18 of Reg. (EC) No 1272/2008 (CLP): PAS[®]; Triclopyr-2-butoxyethyl ester

WARNING

May cause an allergic skin reaction. May cause damage to kidneys through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects

Do not breathe mist/vapours/spray. Wear protective gloves/protective clothing.

IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/attention. Collect spillage.

Dispose of contents/container to a licensed waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT INFORMATION.

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crop/Situation: Grassland

Maximum Individual Dose: 2.0 litres product per hectare

Maximum Total Dose: 2.0 litres product/hectare/annum

Latest Time of Application: 7 days before grazing or harvest


Other Specific Restrictions:

Livestock must be kept out of treated areas for at least 7 days following treatment. IF RAGWORT IS PRESENT, FOLLOW THE GUIDANCE IN THE 'DIRECTIONS FOR USE'.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.



9 990720791612



DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

NOTES

Grass and weeds must be actively growing to ensure good weed control and minimal check to the grass.

Do not spray in drought, very hot or very cold weather.

Control may be reduced if rain falls within 2 hours of application.

Do not roll or harrow grass for 10 days before or 7 days after application.

Clover will be killed or severely checked by application of PAS.

DO NOT sow kale, swedes, turnips or grass mixtures containing clover by direct drilling or minimum cultivation techniques within 6 weeks of applying PAS.

PAS is safe to established grassland. Very occasionally some yellowing of the sward may occur; this is transient and quickly outgrown.

Occasionally, e.g. in hot conditions, vapour drift may occur, making it particularly important to assess the risk to neighbouring vegetation.

Take care to avoid drift onto susceptible crops, non-target plants or waterways. Do not apply directly to, or allow spray drift to come into contact with agricultural or horticultural crops, amenity plantings, gardens, ponds, lakes or watercourses.

Wash out spray equipment thoroughly with water and detergent immediately after use. Traces of PAS could cause harm to susceptible crops sprayed later.

DO NOT use on crops intended for seed production.

Treated crops should not be re-entered before spray deposits on leaf surfaces have completely dried.

Grazing Interval

Exclude livestock during treatment and do not allow livestock to graze treated grassland for at least 7 days following treatment. Where ragwort is present users should consult the Code of Practice on How to Prevent the Spread of Ragwort. Ragwort plants sprayed with this herbicide are more palatable with higher levels of toxins. Animals should be excluded from treated areas until any ragwort has completely recovered or died and there is no visual sign of the dead weed. Do not include treated ragwort in hay or silage crops.

GENERAL INFORMATION

PAS is a foliar acting herbicide for the control of broad-leaved dock and curled dock in established grassland.

ESTABLISHED GRASSLAND

WEEDS CONTROLLED

PAS is recommended for the control of broad-leaved dock and curled dock, in the rosette stage up to 200 mm high or across.

Weed	Rate of Use litres product/ha	Optimum timing of application
Curled dock	1.0 + 1.0	At the rosette stage up to 200 mm tall or across
Broad-leaved dock	Spring + Autumn	
Broad-leaved dock	2.0	At the rosette stage up to 200 mm tall or across
Curled dock		

RATE OF USE AND TIMING

Apply PAS at 2.0 litres/ha in 300 to 400 litres of water as an overall treatment in the autumn or spring for the control of docks. Alternatively PAS may be applied in a programme of 1.0 litre/ha in the spring followed by a second application of 1.0 litre/ha in the autumn for the control of docks.

Use the higher volume of water where the weeds are large at the time of application and where the grass is dense.

Application timing of PAS is crucial for good control and must be made when the weeds and grass are actively growing to ensure good weed control and minimal check to the grass. Docks should be treated in the rosette stage and within the size given above.

It is important that there is sufficient leaf area for uptake of the herbicide. This is particularly true on well-established docks which will have an extensive tap root, but which early in the spring or soon after cutting may only have a few small leaves.

If the grass (and hence the weeds) has been cut for conservation hay or silage, or has been grazed, leave for 2 to 3 weeks to allow sufficient regrowth to occur before spraying.

To allow maximum translocation of PAS to the roots do not cut grass for 28 days after application.

On large well-established docks and where there is a large reservoir of seed in the soil a further control programme in the following year may be required.

MIXING

Fill the spray tank half full with water and add the required amount of PAS. Continue filling with water and maintain agitation during filling, while spraying and during any stoppages. Mix up only sufficient spray volume for the area to be treated and use the spray immediately.

SPRAY QUALITY

Apply PAS as a MEDIUM spray as defined by the BCPC system.

Trademark Acknowledgement

PAS is a trademark of the Dow Chemical Company (Dow) or an affiliated company of Dow.

All other brand names used in this document are trademarks of other manufacturers in which proprietary rights may exist.

Dow AgroSciences Conditions of Supply

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



<http://uk.dowagro.com/wp-content/uploads/PAS-SDS.pdf>

Alternatively contact your supplier

SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

- Operators must wear suitable protective clothing (coveralls) and suitable protective gloves when handling the concentrate.
- Operators must wear suitable protective gloves when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin or eyes immediately.

DO NOT BREATHE SPRAY.

WASH HANDS AND EXPOSED SKIN before meals and after work.

DO NOT APPLY via hand held equipment.

Consumer protection:

NOT TO BE USED ON FOOD CROPS.

Environmental protection:

Livestock must be kept out of treated areas for at least 7 days following treatment. IF RAGWORT IS PRESENT, FOLLOW THE GUIDANCE IN THE 'DIRECTIONS FOR USE'.

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with the LERAP requirement.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 metres of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 metre of the top of a ditch which is dry at the time of application. Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with HSE's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

LERAP scheme not applicable in the Republic of Ireland

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

Storage and disposal:

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

KEEP AWAY from food, drink and animal feeding stuffs.

KEEP OUT of reach of children.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.



Product Registration Number: MAPP 17772

An emulsifiable concentrate containing 150 g ae/litre (14.68% w/w) fluroxypyr (present as 216 g/litre of fluroxypyr-meptyl) and 150 g ae/litre (14.68% w/w) triclopyr (present as 209 g/litre triclopyr butoyl).

A foliar acting herbicide for the control of DOCKS in ESTABLISHED GRASSLAND

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.

PROTECT FROM FROST.

2 Litres e

Dow AgroSciences Limited

CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE

Telephone: (01462) 457272 Fax: (01462) 426605

24 Hour Emergency Telephone Number: +44 (0) 1553 761 251

© Trademark of The Dow Chemical Company ("Dow") or an Affiliated Company of Dow

This label is compliant with the CPA Voluntary Initiative Guidance (UK only)



Product Identifier according to Art.18 of Reg. (EC) No 1272/2008 [CLP]: PAS[®]; Triclopyr-2-butoxyethyl ester

WARNING

May cause an allergic skin reaction. May cause damage to kidneys through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects

Do not breathe the mist/vapours/spray. Wear protective gloves/protective clothing.

IF ON SKIN: Wash with plenty of soap and water.

If skin irritation or rash occurs: Get medical advice/attention. Collect spillage.

Dispose of contents/container to a licensed waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT INFORMATION.

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crop/Situation: Grassland

Maximum Individual Dose: 2.0 litres product per hectare

Maximum Total Dose: 2.0 litres product/hectare/annum

Latest Time of Application: 7 days before grazing or harvest

Other Specific Restrictions:

Livestock must be kept out of treated areas for at least 7 days following treatment. IF RAGWORT IS PRESENT, FOLLOW THE GUIDANCE IN THE 'DIRECTIONS FOR USE'.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.



SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

DO NOT APPLY by hand-held equipment.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

AVOID CONTACT WITH EYES.

WASH HANDS AND EXPOSED SKIN before meals and after work.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

IF YOU FEEL UNWELL, seek medical advice (show the label where possible).

Environmental protection:

Livestock must be kept out of treated areas for at least 7 days following treatment. IF RAGWORT IS PRESENT, FOLLOW THE GUIDANCE IN THE 'DIRECTIONS FOR USE'.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from yards and roads.

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top

of a ditch which is dry at the time of application. Aim spray away from water. This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years. LERAP scheme not applicable in the Republic of Ireland.


Storage and disposal:

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.



Product Registration Number: MAPP 17777

A soluble concentrate containing 200 g/litre (17.4% w/w) clopyralid (present as 305 g/litre of clopyralid as the triethylammonium salt) and 200 g/litre (17.4% w/w) triclopyr (present as 279 g/litre triclopyr as the triethylammonium salt).

A foliar acting herbicide for the control of CREEPING THISTLE in ESTABLISHED GRASSLAND.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work (UK only).

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.

PROTECT FROM FROST.

2 Litres e


Dow AgroSciences Limited

CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE

Telephone: (01462) 457272 Fax: (01462) 426605

24 Hour Emergency Telephone Number: +44 (0) 1553 761251

®Trademark of the Dow Chemical Company ("Dow") or an affiliated company of Dow



Product Identifier according to Art.18 of Reg. (EC) No 1272/2008 [CLP]:Tor®; Clopyralid Triethylamine Salt; Triclopyr Triethylamine Salt; Triethylamine

Danger

Causes serious eye damage.

May cause respiratory irritation.

May cause damage to kidneys through prolonged or repeated exposure.

Very Toxic to aquatic life with long lasting effects.

Do not breathe mist/vapours/spray.

Wear eye protection/face protection.

IF INHALED: Remove person to fresh air and keep comfortable for breathing.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Immediately call a POISON CENTRE/doctor.

Dispose of contents/container to a licensed waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT INFORMATION.

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops/Situations:	Permanent grassland, rotational grass
Maximum Individual Dose:	1.0 litre product per hectare
Maximum Number of Treatments:	1.0 litre product/hectare/annum
Latest Time of Application:	7 days before grazing or harvest - cutting for hay and silage

Other specific restriction

Applications must not be made outside of the period 1st March to 31st October. To protect groundwater do not apply to leys less than 1 year old.

Livestock must be kept out of the treated areas for at least 7 days following treatment. IF RAGWORT IS PRESENT, FOLLOW THE GUIDANCE IN THE 'DIRECTIONS FOR USE' **READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**



DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product. TOR® is a foliar acting herbicide for the control of creeping thistle in established grassland.

NOTES

Grass and weeds must be actively growing to ensure good weed control and minimal check to the grass.

Do not spray in drought, very hot or very cold weather.

Control may be reduced if rain falls within 2 hours of application.

Do not roll or harrow grass for 10 days before or 7 days after application.

Clover will be killed or severely checked by application of TOR.

DO NOT sow kale, swedes, turnips or grass mixtures containing clover by direct drilling or minimum cultivation techniques within 6 weeks of applying TOR. TOR residues in plant tissues which have not completely decayed may affect succeeding susceptible crops e.g. peas, beans and other legumes, carrots and umbelliferae, potatoes and tomatoes, lettuce and other compositae, glasshouse and protected crops. Do not plant susceptible autumn sown crops e.g. winter beans in the same year as treatment with TOR. Where susceptible crops are to be planted in the spring, do not apply TOR later than the end of July of the previous year. Following good agronomic practice ensure that plant remains have completely decayed before planting susceptible crops.

Do not use any plant material treated with TOR for composting or mulching.

Do not use manure for composting from animals fed on crops treated with TOR.

TOR is safe to grass. Very occasionally some yellowing of the sward may occur; this is transient and quickly outgrown.

Occasionally, e.g. in hot conditions, vapour drift may occur, making it particularly important to assess the risk to neighbouring vegetation.

Take extreme care to avoid drift onto crops and non-target plants outside the target area.

All conifers, especially pine and larch, are very sensitive to spray drift.

Grazing Interval

Exclude livestock during treatment and do not allow livestock to graze treated grassland for at least 7 days following treatment. Where ragwort is present users should consult the Code of Practice on How to Prevent the Spread of Ragwort. Ragwort plants sprayed with this herbicide are more palatable with higher levels of toxins. Animals should be excluded from treated areas until any ragwort has completely recovered or died and there is no visual sign of the dead weed. Do not include treated ragwort in hay or silage crops.

WEEDS CONTROLLED

TOR is recommended for the control of creeping thistle in the rosette stage up to 200 mm high or across in permanent grassland and rotational grass.

RATE OF USE

Apply TOR at 1.0 litre/ha in 200 to 400 litres of water as an overall treatment for the control of creeping thistle. Use the higher volume of water where the weeds are large at the time of application and where grass is dense.

TIMING

Application timing of TOR is crucial for good control and must be made when the weeds and grass are actively growing to ensure good weed control and minimal check to the grass. Creeping thistle should be treated in the rosette stage, before stem extension, and within the size given above.

It is important that there is sufficient leaf area for uptake of the herbicide especially soon after cutting where the plant may only have a few small leaves. This is particularly true on well-established thistles, which will have extensive root systems that will generate many new shoots.

If the grass (and hence the weeds) has been cut for conservation hay or silage, or has been grazed, leave for 2 to 3 weeks to allow sufficient regrowth to occur before spraying.

To allow maximum translocation of TOR to the roots do not cut grass for 28 days after application.

On large well-established thistles and where there is a large reservoir of seed in the soil a further application in the following year may be required.

MIXING

Fill the spray tank half full with water and add the required amount of TOR. Continue filling with water and maintain agitation during filling, while spraying and during any stoppages. Mix up only sufficient spray volume for the area to be treated and use the spray immediately.

Wash out spray equipment thoroughly with water and detergent immediately after use. Spray out, fill with clean water and leave overnight. Spray out again before storing or using another product. Traces of TOR could cause harm to susceptible crops sprayed later.

SPRAY QUALITY

Apply TOR as a MEDIUM spray as defined by the BCPC system.

Dow AgroSciences Conditions of Supply

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.



To access the Safety Data Sheet for this product scan the QR code or use the weblink below:

<http://uk.dowagro.com/wp-content/uploads/TOR-SDS.pdf>

Alternatively contact your supplier

SPECIMEN

**THIS PAGE HAS BEEN LEFT
BLANK INTERTIONALLY**

SPECIMEN

SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

DO NOT APPLY by hand-held equipment.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

AVOID CONTACT WITH EYES.

WASH HANDS AND EXPOSED SKIN before meals and after work.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

IF YOU FEEL UNWELL, seek medical advice (show the label where possible).

Environmental protection:

Livestock must be kept out of treated areas for at least 7 days following treatment. IF RAGWORT IS PRESENT, FOLLOW THE GUIDANCE IN THE 'DIRECTIONS FOR USE'.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from yards and roads.

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top

of a ditch which is dry at the time of application. Aim spray away from water. This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years. LERAP scheme not applicable in the Republic of Ireland.

Storage and disposal:

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

 Dow AgroSciences

Tor[®]

HERBICIDE

Product Registration Number: MAPP 17777

A soluble concentrate containing 200 g/litre (17.4% w/w) clopyralid (present as 305 g/litre of clopyralid as the triethylammonium salt) and 200 g/litre (17.4% w/w) triclopyr (present as 279 g/litre triclopyr as the triethylammonium salt).

A foliar acting herbicide for the control of CREEPING THISTLE in ESTABLISHED GRASSLAND.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work (UK Only).

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.

PROTECT FROM FROST.

2 Litres e


Dow AgroSciences Limited

CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE

Telephone: (01462) 457272 Fax: (01462) 426605

24 Hour Emergency Telephone Number: +44 (0) 1553 761251

©Trademark of the Dow Chemical Company ("Dow") or an affiliated company of Dow



Product Identifier according to Art.18 of Reg. (EC) No 1272/2008 [CLP]:Tor[®]; Clopyralid Triethylamine Salt; Triclopyr Triethylamine Salt; Triethylamine

Danger

Causes serious eye damage.

May cause respiratory irritation.

May cause damage to kidneys through prolonged or repeated exposure.

Very Toxic to aquatic life with long lasting effects.

Do not breathe mist/vapours/spray.

Wear eye protection/face protection.

IF INHALED: Remove person to fresh air and keep comfortable for breathing.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Immediately call a POISON CENTRE/doctor.

Dispose of contents/container to a licensed waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT INFORMATION.

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops/Situations:

Permanent grassland, rotational grass

Maximum Individual Dose:

1.0 litre product per hectare

Maximum Number of Treatments:

1.0 litre product/hectare/annum

Latest Time of Application:

7 days before grazing or harvest - cutting for hay and silage

Other specific restriction

Applications must not be made outside of the period 1st March to 31st October. To protect groundwater do not apply to leys less than 1 year old.

Livestock must be kept out of the treated areas for at least 7 days following treatment. IF RAGWORT IS PRESENT, FOLLOW THE GUIDANCE IN THE 'DIRECTIONS FOR USE'

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.