

**GENERAL INFORMATION:**

INTERFACE® is a broad spectrum foliar fungicide exhibiting both contact and mesostemic mode of action. It is a combination of dicarboxamide and strobilurin fungicides with preventative, curative and eradicated properties for use on a range of diseases on managed amenity turf. INTERFACE® acts against all stages of the disease cycle from the inhibition of spore germination to mycelial growth and sporulation.

**RESTRICTIONS:**

INTERFACE® is not to be used on commercial, residential or estate lawns.

Do not apply during drought conditions or to frozen turf.

If grass is being mown, INTERFACE® should be applied after cutting and further mowing should not take place until AT LEAST 48 hours after treatment.

A minimum of four weeks must be observed between treatments.

Do not contaminate water or ditches with the product or its container.

Avoid contamination via drains from farmyards and roads.

Do not clean application equipment near surface water.

A minimum of four weeks must be observed between treatments.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

**DIRECTIONS FOR USE:**

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

**APPLICATION:**

For control of Fusarium Patch, Red Thread and Dollar Spot. For control of Leaf Spot and Rust and for moderate control of Anthracnose where present at time of application against other listed diseases. Use 10 L of INTERFACE® in 400-600 litres of water per hectare of turf. For use with knapsack sprayers or vehicle mounted hydraulic sprayers.

Application Equipment			
Vehicle-mounted Hydraulic Sprayer		Knapsack Sprayer	
Rate per hectare	Water per hectare	Rate per 100 m <sup>2</sup>	Water per 100 m <sup>2</sup>
10 Litres	400-600 litres	100 mls	4 -6 litres
		Rate per 500 m <sup>2</sup>	Water per 500 m <sup>2</sup>
		500 mls	20-30 litres

**Application Timing:**

INTERFACE® can be used when grass is either dormant or actively growing and displays protective, curative and eradicated activity. For best results apply preventatively or at first sign of disease and repeat as necessary at intervals of 28 days or greater.

INTERFACE® can be applied year round.

**Resistance Management:**

INTERFACE® contains active ingredients from strobilurin class of chemistry (Trifloxystrobin) and the dicarboxamide class of chemistry (Iprodione).

The use of INTERFACE® should conform to resistance management strategies. Such strategies should include using in rotation with products containing an active substance with a different mode of action which is effective against the target disease, limiting the total number of applications per season, and using non-chemical methods of disease control. Bayer recommends responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

Maximum number of treatments: 4 per year.

This strategy should include using in rotation with products that have a different mode of action or different chemical group that are effective against the target disease.

**Mixing:**

Sprayers should be THOROUGHLY CLEANED before use and filters and jets checked for damage and blockages.

Thoroughly shake the pack before use.

Half fill the sprayer tank with clean water and agitate. Add the required amount of INTERFACE®.

Mix thoroughly and make up to the required volume of water while still agitating.

Spray immediately after mixing. Continue agitation at all times during spraying and stoppages, until the tank is completely empty.

Wash out all spray equipment thoroughly with water immediately after use.

**SAFETY PRECAUTIONS****Operator Protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment :

Wear suitable protective gloves when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

To protect aquatic organisms respect an unsprayed buffer zone distance to surface water bodies in line with LERAP requirements.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

**Environmental Protection**

DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with chemical or used container.

A 5 m buffer zone has to be applied when standard application equipment is used; DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application.

DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water.

DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water.

DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water.

DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water.

DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water.


DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water.

DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water.

DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water.

DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water.

DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water.



# Interface®

**FUNGICIDE**

- ✓ 1ST UK FUNGICIDE WITH STRESSGARD™ FORMULATION TECHNOLOGY
- ✓ PREVENTATIVE, CURATIVE AND ERADICANT CONTROL

INTERFACE® is a broad spectrum contact and mesostemic foliar fungicide.

For the control of Fusarium Patch (*Microdochium nivale*), Red Thread (*Laetisaria fuciformis*) and Dollar Spot (*Sclerotinia homoeocarpa*).

For control of Leafspot (*Drechslera* spp.), and Rust (*Puccinia* spp) and moderate control of Anthracnose (*Colletotrichum cereale*), where present at the time of application against other listed diseases

MAPP Number 16060

INTERFACE® is an aqueous suspension concentrate containing 256 g/L iprodione and 16 g/L trifloxystrobin

# 5 L e

**Important Information:**

A minimum of four weeks must be observed between treatments.

FOR USE ONLY AS A HORTICULTURAL FUNGICIDE

Situation: Managed amenity turf.

Maximum individual dose : 10 L/ha.

Maximum number of treatments : 4 per year.

Reapplication interval : 28 days.

(i) INTERFACE® may be applied to managed amenity turf sites including:

(a) bowling greens, cricket pitches (excluding outfield), lawn tennis courts and professional sports grounds/stadiums.



(b) golf greens, tees and patch treatment of up to 5% of golf course fairways.

(ii) A minimum of four weeks must be observed between treatments.

(iii) This product must not be used on commercial, residential or estate lawns.

**Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Follow the code of practice for using plant protection products.**

**WARNING**

Suspected of causing cancer.  
Very toxic to aquatic life with long lasting effects.  
Wear protective gloves/protective clothing/eye protection/face protection.  
IF exposed or concerned: Get medical advice/attention.  
Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.  
To avoid risks to human health and the environment, comply with the instructions for use.  
Contains trifloxystrobin. May produce an allergic skin reaction.

**MAPP: 16060**

GB80917031C - ARTICLE 80948433 - RA3

For 24 hour emergency information contact Bayer CropScience Limited  
Telephone: 0800 220876

Batch number : See on the pack

# Bayer

Bayer CropScience Limited  
230 Cambridge Science Park  
Milton Road  
Cambridge CB4 0WB UK  
Tel: 00800 12149451  
www.escience.bayercropscience.com  
for MSDS and larger label

Interface® is a registered trademark of Bayer CropScience Ltd